

Introduction to Research Integrity and Research Ethics and Responsible Conduct of Research: From a Veterinarian's Perspective

By

Nino Arias, *MV Dr.med.vet.*

January 30th, 2015

Conducta Responsable en Investigación
Curso de autoaprendizaje

<http://goo.gl/WYcr8u>

To my personal perspective :

Assure good welfare for animals
participating in research experiments

Physiological measures of
good or poor animal welfare

My choice:
Neuroendocrine system

Faecal cortisol metabolites in South American camelids

- After ACTH challenge, plasma cortisol and FCM levels were increased in alpacas and llamas.
- The persecution, capture, and confinement increased FCM values in llamas and vicuñas.
- FCM 11,17-DOA are suited for **non-invasive** measurement of adrenocortical activity in alpacas, llamas and vicuñas.

References

- Arias N, Requena M, Palme R. 2013. Measuring faecal glucocorticoid metabolites as a noninvasive tool for monitoring adrenocortical activity in South American camelids. *Anim Welfare* 22: 25-31.
- Bravo PW, Garnica J, Aviles E. 2001. Cortisol concentrations in the perinatal and weaning periods of alpacas. *Anim Reprod Sci* 67: 125-129.
- Carmanchahi PD, Ovejero R, Marull C, López GC, Schroeder N, Jahn GA, et al. 2011. Physiological response of wild guanacos to capture for live shearing. *Wildlife Res* 38: 61-68.
- Concepts in Animal Welfare. 2007. An animal welfare syllabus. University of Bristol. World Society for the Protection of Animals WSPA.
- Möstl E and Palme R. 2002. Hormones as indicators of stress. *Domestic Animal Endocrinology* 23: 67-74.
- Möstl E, Rettenbacher S, Palme R. 2005. Measurement of corticosterone metabolites in birds' droppings: An analytical approach. *Annals New York Acad. Sci.* 1046, 17-34.
- Palme R. 2012. Monitoring stress hormone metabolites as a useful, non-invasive tool for welfare assessment in farm animals. *Anim Welfare* 21: 331-337.